

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	CALCR
Synonyms	CRT; CT-R; CTR; CTR1
Description	Human CALCR full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P30988
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Neuroactive ligand-receptor interaction
Molecular Weight	The human full length CALCR protein has a MW of 55.3 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein is a high affinity receptor for the peptide hormone calcitonin and belongs to a subfamily of seven transmembrane-spanning G protein-coupled receptors. The encoded protein is involved in maintaining calcium homeostasis and in regulating osteoclast-mediated bone resorption. Polymorphisms in this gene have been associated with variations in bone mineral density and onset of osteoporosis.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate CALCR-Nanodisc 0.2 μ g Human CALCR-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag CALCR-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CALCR-Nanodisc is 0.706ng/ml.




Figure 2. Human CALCR-Nanodisc, Flag Tag on SDS-PAGE

