

PRODUCT INFORMATION

C-Flag Tag Tag

CAV1 **Target**

Synonyms BSCL3; CGL3; LCCNS; MSTP085; PPH3; VIP21

Description Human CAV1 full length protein-MNP

Delivery In Stock **Uniprot ID** Q03135 **Expression Host HEK293**

Storage & Shipping

Background

Druggable Genome, Transmembrane **Protein Families**

Protein Pathways Focal adhesion, Viral myocarditis

The human full length CAV1 protein has a MW of **Molecular Weight**

20.3 kDa

Lyophilized from PBS. Normally 5% - 8% Formulation & trehalose is added as protectants before

Reconstitution lyophilization. Please see Certificate of Analysis

for specific instructions. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

The scaffolding protein is the main component of the caveolae plasma membranes found in most cell types. The protein links integrin subunits to the tyrosine kinase FYN, an initiating step in coupling integrins to the Ras-ERK pathway and promoting cell cycle progression. The gene is a tumor suppressor gene candidate and a negative regulator of the Ras-p42/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are

located next to each other on chromosome 7 and express colocalizing proteins that form a stable hetero-oligomeric complex. Mutations in this gene have been associated with Berardinelli-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of

caveolin 1.

Usage Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)







ELISA assay to evaluate CAV1-MNP 0.5µg Human CAV1-MNP per well




Figure 1. Elisa plates were pre-coated with $0.5\mu g/per$ well purified human CAV1 full length membrane nanoparticles. Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CAV1 full length membrane nanoparticles is 5.552ng/ml.

Email: info@dimabio.com Website: www.dimabio.com

