

PRODUCT INFORMATION

TargetCAV1SynonymsBSCL3; CGL3; LCCNS; MSTP085; PPH3; VIP21DescriptionHuman CAV1-Strep full length protein-synthetic nanodiscDelivery6-8wecksUniprot IDQ03135Expression HostHEK293Protein FamiliesDruggable Genome, TransmembraneProtein PathwaysFocal adhesion, Viral myocarditisMolecular WeightThe human full length CAV1-Strep protein has a MW of 20.3 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCI, pH 8.0). Normally 5% or 8% trehalose is added as protectants before lyophilized form. Please see Certificate of Analysis forStorage & ShippingStore at -20°C to -80°C for 12 months in lyophilized form. The reconstitution, if not intended for use within a month, aliquot and store at -80°C to before glyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C in the reconstitution, if not intended for use within the main component of the caveolae plasma membranes found in most store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a daveolin 2 are loculation of the casedolate and a negative eregulator of the Ras-P42/44 mitogen-activated kinase exacted. Caveolin 1 and caveolin 2 are located next to each other on chromosome 7 and express colocalizing protein stat form a stable heave been associated with Beardinelli-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of caveolin 1.BackgroundResearch use o	Тад	C-Flag&Strep Tag
DescriptionHuman CAV1-Strep full length protein-synthetic nanodiscDelivery6~8weeksUniprot IDQ03135Expression HostHEK293Protein FamiliesDruggable Genome, TransmembraneProtein PathwaysFocal adhesion, Viral myocarditisMolecular WeightThe human full length CAV1-Strep protein has a MW of 20.3 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCI, pH 8.0). Normally 5% - 8% trehalose is added as protectants before yophilization. Please see Certificate of Analysis forStorage & ShippingStorage & ShippingBackgroundHe scaffolding protein is the main component of the caveolae plasm amembranes found in most cell types. The protein links integrin subunits to the typosine kinase FXN, an initiating step in coupling integrins to the Ras-ERK pathway and promoting cell cycle progression. The gene is a turos to suppressor gene candidate and a negative regulator of the Ras-PL/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are located next to each other on chromosome 7 and express colocalizing protein stat form a stable hetero-oligomeric complex. Mutations in this gene have been associated with Berardinelli-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of caveolin 1.	Target	CAV1
DescriptionnanodiscDelivery6-8weeksUniprot IDQ03135Expression HostHEK293Protein FamiliesDruggable Genome, TransmembraneProtein FamiliesDruggable Genome, TransmembraneProtein PathwaysFocal adhesion, Viral myocarditisMolecular WeightThe human full length CAV1-Strep protein has a MW of 20.3 kDa Lyophilization Duffer (20 mM Tris-HCI, 10 nm NACI, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis forStorage & ShippingStore at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. The scaffolding protein is the main component of the eaveloae plasma membranes found in most ceell types. The protein links integrin subunits to the tyrosine kinase FYN, an initiating step in coupling integrins to the Ras-ERK pathway and promoting cell cycle progression. The gene is a tumor suppressor gene candidate and a negative regulator of the Ras-P42/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are located next to each other on chromosome 7 and express colocalizing proteins that form a stable hetero-oligomeric complex. Mutations in this gene have been associated with Berardinell-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of caveolin 1.UsageResearch use only	Synonyms	BSCL3; CGL3; LCCNS; MSTP085; PPH3; VIP21
Uniprot IDQ03135Expression HostHEK293Protein FamiliesDruggable Genome, TransmembraneProtein PathwaysFocal adhesion, Viral myocarditisMolecular WeightThe human full length CAV1-Strep protein has a MW of 20.3 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilizet of contact and the second stress of the second stress o	Description	
Expression HostHEK293Protein FamiliesDruggable Genome, TransmembraneProtein PathwaysFocal adhesion, Viral myocarditisMolecular WeightThe human full length CAV1-Strep protein has a MW of 20.3 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilized for use within a month, aliquot and store at -20°C to -80°C for 12 months in lyophilized proteins are shipped at ambient temperature.Storage & ShippingThe scaffolding protein is the main component of the caveolae plasma membranes found in most cell types. The protein 1 mis integrine suburits to the tyrosine kinase FYN, an initiating step in coupling integrins to the Ras-FEK pathway and promoting cell cycle progression. The gene is a tumor suppressor gene candidate and a nagative regulator of the Ras-PEK/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are looded next to each other on chromsome 7 and express colocalizing proteins that form a stable hetero-oligomeric complex. Mutations in this gene have been associated with Berardinelli-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of caveolin 1.UsageResearch use only	Delivery	6~8weeks
Protein FamiliesDruggable Genome, TransmembraneProtein PathwaysFocal adhesion, Viral myocarditisMolecular WeightThe human full length CAV1-Strep protein has a MW of 20.3 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HC1, 150 mM NaCl, pH 8.00. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis forStorage & ShippingStore at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.BackgroundBackgroundThe scaffolding protein is the main component of the caveolae plasma membranes found in most cell types. The protein links integrin subunits to the twrosine kinase FYN, an initiating step in coupling integrins to the Ras-P42/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are located next to each other on chromosome 7 and express colocalizing proteins that form a stable hetero-oligomeric complex. Mutations in this gene have been associated with Berardinelli-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of caveolin 1.UsageResearch use only	Uniprot ID	Q03135
Protein PathwaysFocal adhesion, Viral myocarditisMolecular WeightFocal adhesion, Viral myocarditisMolecular WeightThe human full length CAV1-Strep protein has a MW of 20.3 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCL 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis forStorage & ShippingStore at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.BackgroundThe scaffolding protein is the main component of the caveolae plasma membranes found in most cell types. The protein links integrin subunits to the tyrosine kinase FYN, an initiating step in coupling integrins to the Ras-ERK pathway and promoting cell cycle progression. The gene is a tumor suppressor gene candidate and a negative regulator of the Ras-p42/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are located next to each other on chromosome 7 and express colocalizing proteins that form a stable hetero-oligomeric complex. Mutations in this gene have been associated with Berardinelli-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of caveolin 1.WageResearch use only	Expression Host	HEK293
Molecular WeightThe human full length CAV1-Strep protein has a MW of 20.3 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis forStorage & ShippingStorage & ShippingStore at -20°C to -80°C for 12 months in lyophilized from. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. The scaffolding protein is the main component of the caveolae plasma membranes found in most cell types. The protein links integrin subunits to the tyrosine kinase FN, an initiating step in coupling integrins to the Ras-ERK pathway and promoting cell cycle progression. The gene is a tumor suppressor gene candidate and a negative regulator of the Ras-p42/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are located next to each other on chromosome 7 and express colocalizing proteins that form a stable hetero-oiligomeric complex. Mutations in this gene have been associated with Berardinelli-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of caveolin 1.UsageResearch use only	Protein Families	Druggable Genome, Transmembrane
Molecular WeightMW of 20.3 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis forStorage & ShippingStore at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.BackgroundThe scaffolding protein is the main component of the tyrosine kinase FYN, an initiating step in coupling integrins to the Ras-ERK pathway and promoting cell cycle progression. The gene is a tumor suppressor gene candidate and a negative regulator of the Ras-p42/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are located next to each other on chromosome 7 and express colocalizing proteins that form a stable hetero-oligomeric complex. Mutations in this gene have been associated with Berardineli-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of caveolin 1.UsageResearch use only	Protein Pathways	Focal adhesion, Viral myocarditis
Formulation & ReconstitutionLyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis forStorage & ShippingStore at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.BackgroundThe scaffolding protein is the main component of the caveolae plasma membranes found in most cell types. The protein links integrin subunits to the tyrosine kinase FYN, an initiating step in coupling integrins to the Ras-ERK pathway and promoting cell cycle progression. The gene is a tumor suppressor gene candidate and a negative regulator of the Ras-p42/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are located next to each other on chromosome 7 and express colocalizing proteins that form a stable hetero-oligomeric complex. Mutations in this gene have been associated with Berardinelli-Seip orgenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of caveolin 1.UsageResearch use only	Molecular Weight	
Storage & ShippingIvophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.BackgroundThe scaffolding protein is the main component of the caveolae plasma membranes found in most cell types. The protein links integrin subunits to the tyrosine kinase FYN, an initiating step in coupling integrins to the Ras-ERK pathway and promoting cell cycle progression. The gene is a tumor suppressor gene candidate and a negative regulator of the Ras-p42/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are located next to each other on chromosome 7 and express colocalizing proteins that form a stable hetero-oligomeric complex. Mutations in this gene have been associated with Berardinelli-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of caveolin 1.UsageResearch use only		Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis
 Background Background The scaffolding protein is the main component of the caveolae plasma membranes found in most cell types. The protein links integrin subunits to the tyrosine kinase FYN, an initiating step in coupling integrins to the Ras-ERK pathway and promoting cell cycle progression. The gene is a tumor suppressor gene candidate and a negative regulator of the Ras-p42/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are located next to each other on chromosome 7 and express colocalizing proteins that form a stable hetero-oligomeric complex. Mutations in this gene have been associated with Berardinelli-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of caveolin 1. Usage 	Storage & Shipping	lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient
	Background	the caveolae plasma membranes found in most cell types. The protein links integrin subunits to the tyrosine kinase FYN, an initiating step in coupling integrins to the Ras-ERK pathway and promoting cell cycle progression. The gene is a tumor suppressor gene candidate and a negative regulator of the Ras-p42/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are located next to each other on chromosome 7 and express colocalizing proteins that form a stable hetero-oligomeric complex. Mutations in this gene have been associated with Berardinelli-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of
	Usage	Research use only
Conjugate Unconjugated	Conjugate	Unconjugated

Email: info@dimabio.com Website: www.dimabio.com

