

## **PRODUCT INFORMATION**

Tag C-Flag Tag

CB2 **Target** 

**Synonyms** CB-2; CNR2; CX5

**Description** Human CB2 full length protein-synthetic nanodisc

**Delivery** In Stock **Uniprot ID** P34972 **Expression Host HEK293** 

**Protein Families** Druggable Genome, GPCR, Transmembrane

**Protein Pathways** Neuroactive ligand-receptor interaction

The human full length CB2 protein has a MW of Molecular Weight

39.7 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% Formulation & - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The cannabinoid delta-9-tetrahydrocannabinol is the principal psychoactive ingredient of

marijuana. The proteins encoded by this gene and the cannabinoid receptor 1 (brain) (CNR1) gene have the characteristics of a guanine nucleotidebinding protein (G-protein)-coupled receptor for cannabinoids. They inhibit adenylate cyclase activity in a dose-dependent, stereoselective, and

pertussis toxin-sensitive manner. These proteins have been found to be involved in the cannabinoid-induced CNS effects (including alterations in mood and cognition) experienced by users of marijuana. The cannabinoid receptors are members of family 1 of the G-protein-coupled

receptors.


Usage Research use only

Conjugate Unconjugated

**Background** 



Email: info@dimabio.com Website: www.dimabio.com







## ELISA assay to evaluate CB2-Nanodisc 0.2µg Human CB2-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag CB2-Nanodisc ( $0.2\mu g/per$  well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CB2-Nanodisc is 1.557ng/ml.




Figure 2. Human CB2-Nanodisc, Flag Tag on SDS-PAGE

Email: info@dimabio.com Website: www.dimabio.com

