

PRODUCT INFORMATION

C-Flag Tag Tag CACNG1 **Target CACNLG Synonyms**

Human CACNG1 full length protein-synthetic **Description**

nanodisc **Delivery** In Stock **Uniprot ID** Q06432 **Expression Host HEK293**

Storage & Shipping

Background

Druggable Genome, Ion Channels: Other, **Protein Families**

Transmembrane

Arrhythmogenic right ventricular cardiomyopathy (ARVC), Cardiac muscle contraction, Dilated

Protein Pathways cardiomyopathy, Hypertrophic cardiomyopathy (HCM), MAPK signaling pathway

The human full length CACNG1 protein has a MW **Molecular Weight**

of 25.0 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% Formulation & 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

Voltage-dependent calcium channels are composed of five subunits. The protein encoded by this gene represents one of these subunits, gamma, and is one of two known gamma subunit proteins. This particular gamma subunit is part of skeletal muscle 1,4-dihydropyridine-sensitive calcium changes and integral membrane protein that plays a role in excitation-contraction

coupling. This gene is part of a functionally diverse eight-member protein subfamily of the PMP-22/EMP/MP20 family and is located in a cluster with two family members that function as

transmembrane AMPÁ receptor regulatory proteins (TARPs).

Usage Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com



ELISA assay to evaluate CACNG1-Nanodisc 0.2μg Human CACNG1-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag CACNG1-Nanodisc ($0.2\mu g/per$ well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CACNG1-Nanodisc is 4.936ng/ml.




Figure 2. Human CACNG1-Nanodisc, Flag Tag on SDS-PAGE

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

