## **PRODUCT INFORMATION**

**Target** CCL2

**Synonyms** HC11;MCAF;MCP-1

Recombinant human CCL2 protein with C-terminal Description

human Fc tag

**Delivery** In Stock **Uniprot ID** P13500 **Expression Host HEK293** 

Tag C-Human Fc Tag

Molecular

Storage & Shipping

**Background** 

CCL2(Gln24-Thr99) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of

34.8 kDa after removal of the signal peptide. The **Molecular Weight** apparent molecular mass of CCL2-hFc is

approximately 35-55 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Chemokines are a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. This chemokine is a member of the CC subfamily which is characterized by two

adjacent cysteine residues. This cytokine displays chemotactic activity for monocytes and basophils but not for neutrophils or eosinophils. It has been


implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis and

atherosclerosis. It binds to chemokine receptors CCR2 and CCR4. Elevated expression of the encoded protein is associated with severe acute

> Email: info@dimabio.com Website: www.dimabio.com

respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. [provided by RefSeq, Aug 2020]

Usage Research use only Conjugate Unconjugated








Figure 1. Human CCL2 Protein, hFc Tag on SDS-PAGE under reducing condition.

## Human CCL2, hFc tagged protein ELISA

0.2 μg of Human CCL2, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human CCL2 Protein, hFc Tag (PME100919) can bind Anti-CCL2 Neutralizing antibody BME100125 in a linear range of 0.92–234.38 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

