

PRODUCT INFORMATION

Tag C-Flag Tag

CCR3 **Target**

CC-CKR-3; C C CKR3; CD193; CKR 3; CKR3; **Synonyms**

CMKBR3

Description Human CCR3 full length protein-MNP

Delivery In Stock **Uniprot ID** P51677 **Expression Host HEK293**

Storage & Shipping

Background

Protein Families Druggable Genome, GPCR, Transmembrane

Chemokine signaling pathway, Cytokine-cytokine **Protein Pathways**

receptor interaction

The human full length CCR3 protein has a MW of **Molecular Weight**

41.0 kDa

Lyophilized from PBS. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.


The protein is a receptor for C-C type chemokines. It belongs to family 1 of the G protein-coupled receptors. This receptor binds and responds to a variety of chemokines, including eotaxin (CCL11), eotaxin-3 (CCL26), MCP-3 (CCL7), MCP-4 (CCL13), and RANTES (CCL5). It is highly expressed in eosinophils and basophils, and is also detected in TH1 and TH2

cells, as well as in airway epithelial cells. This receptor may contribute to the accumulation and activation of eosinophils and other inflammatory cells in the allergic airway. It is also known to be an entry co-receptor for HIV-1. This gene and seven other chemokine receptor genes form a chemokine receptor gene cluster on the chromosomal region 3p21. Alternatively spliced

transcript variants have been described.

Usage Research use only Conjugate Unconjugated

> Email: info@dimabio.com Website: www.dimabio.com







ELISA assay to evaluate CCR3-MNP 0.5µg Human CCR3-MNP per well





Figure 1. Elisa plates were pre-coated with 0.5µg/per well purified human CCR3 full length membrane nanoparticles. Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CCR3 full length membrane nanoparticles is 23.39ng/ml.



Email: info@dimabio.com Website: www.dimabio.com

