

## **PRODUCT INFORMATION**

Target	CCR3
Synonyms	CC-CKR-3; C C CKR3; CD193; CKR 3; CKR3; CMKBR3
Description	Human CCR3 full length protein-MNP
Delivery	In Stock
Uniprot ID	P51677
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Chemokine signaling pathway, Cytokine-cytokine receptor interaction
Molecular Weight	The human full length CCR3 protein has a MW of 41.0 kDa
Formulation & Reconstitution	Lyophilized from PBS. Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein is a receptor for C-C type chemokines. It belongs to family 1 of the G protein-coupled receptors. This receptor binds and responds to a variety of chemokines, including eotaxin (CCL11), eotaxin-3 (CCL26), MCP-3 (CCL7), MCP-4 (CCL13), and RANTES (CCL5). It is highly expressed in eosinophils and basophils, and is also detected in TH1 and TH2 cells, as well as in airway epithelial cells. This receptor may contribute to the accumulation and activation of eosinophils and other inflammatory cells in the allergic airway. It is also known to be an entry co-receptor for HIV-1. This gene and seven other chemokine receptor genes form a chemokine receptor gene cluster on the chromosomal region 3p21. Alternatively spliced transcript variants have been described.
Usage	Research use only
Conjugate	Unconjugated

Email: info@dimabio.com Website: www.dimabio.com








Figure 1. Elisa plates were pre-coated with 0.5µg/per well purified human CCR3 full length membrane nanoparticles. Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CCR3 full length membrane nanoparticles is 23.39ng/ml.

Email: info@dimabio.com Website: www.dimabio.com

