

PRODUCT INFORMATION

Target CCR3

Synonyms CC-CKR-3; C C CKR3; CD193; CKR 3; CKR3;

CMKBR3

Description Human CCR3 full length protein-MNP

Delivery In Stock
Uniprot ID P51677
Expression Host HEK293

Protein Families Druggable Genome, GPCR, Transmembrane

Protein PathwaysChemokine signaling pathway, Cytokine-cytokine

receptor interaction

Molecular Weight The human full length CCR3 protein has a MW of

41.0 kDa

Formulation &Lyophilized from PBS. Normally 5% – 8% trehalose is added as protectants before

Reconstitution Iyophilization. Please see Certificate of Analysis

for specific instructions.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

Storage & Shipping intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

The protein is a receptor for C-C type chemokines. It belongs to family 1 of the G protein-coupled receptors. This receptor binds and responds to a variety of chemokines, including eotaxin (CCL11), eotaxin-3 (CCL26), MCP-3 (CCL7), MCP-4 (CCL13), and RANTES (CCL5). It is highly expressed in eosinophils and basophils, and is also detected in TH1 and TH2 cells, as well as in airway epithelial cells. This

cells, as well as in airway epithelial cells. This receptor may contribute to the accumulation and activation of eosinophils and other inflammatory cells in the allergic airway. It is also known to be an entry co-receptor for HIV-1. This gene and seven other chemokine receptor genes form a chemokine receptor gene cluster on the chromosomal region 3p21. Alternatively spliced

Email: info@dimabio.com Website: www.dimabio.com

transcript variants have been described.

Usage Research use only

Background

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)



ELISA assay to evaluate CCR3-MNP 0.5µg Human CCR3-MNP per well




Figure 1. Elisa plates were pre-coated with 0.5µg/per well purified human CCR3 full length membrane nanoparticles. Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CCR3 full length membrane nanoparticles is 23.39ng/ml.

