

PRODUCT INFORMATION

C-Flag&Strep Tag Tag

CCR5 **Target**

Expression Host

Storage & Shipping

Background

CC-CKR-5; CCCKR5; CCR-5; CD195; CKR-5; CKR5; **Synonyms**

CMKBR5; IDDM22

Human CCR5-Strep full length protein-synthetic Description

nanodisc

Delivery 6~8weeks **Uniprot ID** P51681 **HFK293**

Druggable Genome, ES Cell Differentiation/IPS. **Protein Families**

GPCR, Transmembrane

Chemokine signaling pathway, Cytokine-cytokine receptor interaction, Endocytosis **Protein Pathways**

The human full length CCR5-Strep protein has a **Molecular Weight**

MW of 40.5 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before Formulation & Reconstitution lyophilization. Please see Certificate of Analysis

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

A member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors.

This protein is expressed by T cells and

macrophages, and is known to be an important co-receptor for macrophage-tropic virus, including HIV, to enter host cells. Defective alleles of this gene have been associated with the HIV infection resistance. The ligands of this receptor include monocyte chemoattractant protein 2 (MCP-2), macrophage inflammatory protein 1 alpha (MIP-1 alpha), macrophage inflammatory protein 1 beta (MIP-1 beta) and regulate so no activation normal T

expressed and secreted protein (RANTES) Expression of this gene was also detected in a promyeloblastic cell line, suggesting that this protein may play a role in granulocyte lineage proliferation and differentiation. This gene is located at the chemokine receptor gene cluster

region. An allelic polymorphism in this gene results in both functional and non-functional alleles; the reference genome represents the

> Email: info@dimabio.com Website: www.dimabio.com

functional allele.

Usage Research use only Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)

