

PRODUCT INFORMATION

Tag C-Flag Tag

CCR7 **Target**

BLR2; CC-CKR-7; CCR-7; CD197; CDw197; **Synonyms**

CMKBR7; EBI1

Human CCR7 full length protein-synthetic Description

nanodisc

Delivery In Stock **Uniprot ID** P32248 **HEK293 Expression Host**

Druggable Genome, GPCR, Transmembrane **Protein Families**

Chemokine signaling pathway, Cytokine-cytokine **Protein Pathways**

receptor interaction

The human full length CCR7 protein has a MW of Molecular Weight

42.9 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% Formulation & 8% trehalose is added as protectants before Reconstitution lyophilization. Please see Certificate of Analysis

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature

The protein is a member of the G protein-coupled receptor family. This receptor was identified as a gene induced by the Epstein-Barr virus (EBV), and is thought to be a mediator of EBV effects on B lymphocytes. This receptor is expressed in various lymphoid tissues and activates B and T lymphocytes. It has been shown to control the

migration of memory T cells to inflamed tissues, as well as stimulate dendritic cell maturation. The chemokine (C-C motif) ligand 19 (CCL19/ECL) has been reported to be a specific ligand of this receptor. Signals mediated by this receptor regulate T cell homeostasis in lymph nodes, and

> Email: info@dimabio.com Website: www.dimabio.com

may also function in the activation and polarization of T cells, and in chronic inflammation pathogenesis.

Usage Research use only Conjugate Unconjugated

Background





ELISA assay to evaluate CCR7-Nanodisc 0.2μg Human CCR7-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag CCR7-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CCR7-Nanodisc is 2.044ng/ml.

Email: info@dimabio.com Website: www.dimabio.com



Figure 2. Human CCR7-Nanodisc, Flag Tag on SDS-PAGE