

## **PRODUCT INFORMATION**

CD155 **Target** 

**Synonyms** PVR;FLJ25946;PVS;CD155;TAGE4;HVED;NECL5

Recombinant human CD155 protein with C-**Description** terminal mouse Fc and 6×His tag

**Delivery** In Stock **Uniprot ID** P15151 **Expression Host HEK293** 

Tag C-Mouse Fc and 6×His Tag

Molecular CD155(Trp21-Asn343) mFc(Pro99-Lys330) 6×His

Characterization

The protein has a predicted molecular mass of 62.2 kDa after removal of the signal peptide. The **Molecular Weight** apparent molecular mass of CD155-mFc-His is

approximately 95-130 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a transmembrane glycoprotein belonging to the immunoglobulin superfamily. The external domain mediates cell attachment to the extracellular matrix molecule vitronectin, while its

intracellular domain interacts with the dynein light chain Tctex-1/DYNLT1. The gene is specific to the primate lineage, and serves as a cellular receptor for poliovirus in the first step of poliovirus replication. Multiple transcript variants


encoding different isoforms have been found for

this gene.

Research use only Usage

Conjugate Unconjugated

**Background** 



/+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com








Figure 1. Human CD155 Protein, mFc-His Tag on SDS-PAGE under reducing condition.

## Human CD155, mFc-His Tagged protein ELISA

0.2 µg of TIGIT, hFc Tagged protein per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human CD155, mFc-His tagged protein (PME100027) can bind Human TIGIT, hFc tagged protein PME100107 in a linear range of 1.95-125 ng/ml.

Email: info@dimabio.com Website: www.dimabio.com





## Human CD155, mFc-His tagged protein ELISA

 $0.1 \mu g$  of Human CD155, mFc-His tagged protein per well




Figure 3. ELISA plate pre-coated by 1  $\mu$ g/mL (100  $\mu$ L/well) Human CD155 Protein, mFc-His Tag(PME100027) can bind Anti-CD155 Rabbit mAb in a linear range of 3.2-80 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

