

## PRODUCT INFORMATION

<b>Tag</b>	C-Flag Tag
<b>Target</b>	CLCN5
<b>Synonyms</b>	CLC5, CLCK2, CIC-5, DENT1, DENTS, NPHL1, NPHL2, XLRH, XRN, hCIC-K2
<b>Description</b>	Human CLCN5 full length protein-synthetic nanodisc
<b>Delivery</b>	6~8weeks
<b>Uniprot ID</b>	P51795
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Ion Channels: Other
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length CLCN5 protein has a MW of 83.1kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a member of the CIC family of chloride ion channels and ion transporters. The encoded protein is primarily localized to endosomal membranes and may function to facilitate albumin uptake by the renal proximal tubule. Mutations in this gene have been found in Dent disease and renal tubular disorders complicated by nephrolithiasis. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jan 2013]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated

