

PRODUCT INFORMATION

| | |
|---|---|
| Target | CLEC4A |
| Synonyms | CD367; CLECSF6; DCIR; DDB27; HDCGC13P; hDCIR; LLIR |
| Description | Recombinant Human CLEC4A Protein with N-terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | Q9UMR7 |
| Expression Host | HEK293 |
| Tag | N-Human Fc Tag |
| Molecular Characterization | hFc(Glu99-Ala330) CLEC4A(Gln70-Leu237) |
| Molecular Weight | The protein has a predicted molecular mass of 46.0 kDa after removal of the signal peptide. The apparent molecular mass of hFc-CLEC4A is approximately 55-70 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Purity | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Formulation & Reconstitution | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Storage&Shipping | This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily. Members of this family share a common protein fold and have diverse functions, such as cell adhesion, cell-cell signalling, glycoprotein turnover, and roles in inflammation and immune response. The encoded type 2 transmembrane protein may play a role in inflammatory and immune response. Multiple transcript variants encoding distinct isoforms have been identified for this gene. This gene is closely linked to other CTL/CTLD superfamily members on chromosome 12p13 in the natural killer gene complex region. [provided by RefSeq, Jul 2008] |
| Background | Research use only |
| Usage | Unconjugated |
| Conjugate | |






Figure 1.Human CLEC4A Protein, hFc Tag on SDS-PAGE under reducing condition.

