

PRODUCT INFORMATION

| | |
|---|--|
| Tag | C-Flag Tag |
| Target | CLTR2 |
| Synonyms | CYSLT2, CYSLT2R, GPCR21, HG57, HPN321, KPG_011, PSEC0146, hGPCR21 |
| Description | Human CLTR2 full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | Q9NS75 |
| Expression Host | HEK293 |
| Protein Families | GPCR, Transmembrane, Druggable Genome, |
| Protein Pathways | GPCRDB Class A Rhodopsin-like, |
| Molecular Weight | The human full length CLTR2 protein has a MW of 39.6kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The cysteinyl leukotrienes LTC4, LTD4, and LTE4 are important mediators of human bronchial asthma. Pharmacologic studies have determined that cysteinyl leukotrienes activate at least 2 receptors, the protein encoded by this gene and CYSLTR1. This encoded receptor is a member of the superfamily of G protein-coupled receptors. It seems to play a major role in endocrine and cardiovascular systems. [provided by RefSeq, Jul 2008] |
| Usage | Research use only |
| Conjugate | Unconjugated |

