

PRODUCT INFORMATION

Target	CS1
Synonyms	SLAMF7;CD319;CS1;CRACC;19A;FOAP-12
Description	Recombinant human CS1 protein with C-terminal human Fc and 6×His tag
Delivery	In Stock
Uniprot ID	Q9NQ25
Expression Host	HEK293
Tag	C-Human Fc and 6×His Tag
Molecular Characterization	CS1(Ser23-Asp223) hFc(Glu99-Ala330) 6×His tag
Molecular Weight	The protein has a predicted molecular mass of 70 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	SLAM family member 7 (SLAMF7) is also known as CD2-like receptor-activating cytotoxic cells (CRACC), Membrane protein FOAP-12, CD antigen CD319, Novel Ly9, Protein 19A, which is a single-pass type I membrane protein and a member of the CD2 family of cell surface receptors. SLAMF7 is expressed in spleen, lymph node, peripheral blood leukocytes, bone marrow, small intestine, stomach, appendix, lung and trachea. Isoform 1 of SLAMF7 mediates NK cell activation through a SH2D1A-independent extracellular signal-regulated ERK-mediated pathway. May play a role in lymphocyte adhesion. Isoform 3 of SLAMF7 does not mediate any NK cell activation.
Usage	Research use only
Conjugate	Unconjugated







Figure 1. Human CS1, hFc-His Tag on SDS-PAGE under reducing condition.

Human CS1, hFc Tagged protein ELISA

0.2 μ g of Human CS1, hFc-His Tagged protein per well

Figure 2. ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human CS1, hFc-His tagged protein (PME100002) can bind Anti-CS1 Neutralizing antibody BME100002 in a linear range of 7.81-210.7 ng/ml.