

## PRODUCT INFORMATION

|   |   |
|---|---|
| <b>Target</b>                           | CXCL1   |
| <b>Synonyms</b>                         | FSP;GRO1;GROa;MGSA;MGSA-a;NAP-3;SCYB1   |
| <b>Description</b>                      | Recombinant Human CXCL1 with N-terminal human Fc tag  |
| <b>Delivery</b>                         | In Stock  |
| <b>Uniprot ID</b>                       | P09341  |
| <b>Expression Host</b>                  | HEK293  |
| <b>Tag</b>                              | N-Human Fc Tag  |
| <b>Molecular Characterization</b>       | hFc(Glu99-Ala330) CXCL1(Ala35-Asn107)   |
| <b>Molecular Weight</b>                 | The protein has a predicted molecular mass of 34.0 kDa after removal of the signal peptide. The apparent molecular mass of hFc-CXCL1 is approximately 35-40 kDa due to glycosylation.   |
| <b>Purity</b>                           | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.  |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Background</b>                       | This antimicrobial gene encodes a member of the CXC subfamily of chemokines. The encoded protein is a secreted growth factor that signals through the G-protein coupled receptor, CXC receptor 2. This protein plays a role in inflammation and as a chemoattractant for neutrophils. Aberrant expression of this protein is associated with the growth and progression of certain tumors. A naturally occurring processed form of this protein has increased chemotactic activity. Alternate splicing results in coding and non-coding variants of this gene. A pseudogene of this gene is found on chromosome 4. [provided by RefSeq, Sep 2014] |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |






Figure 1. Human CXCL1 Protein, hFc Tag on SDS-PAGE under reducing condition.

