

PRODUCT INFORMATION

| | |
|------------------------------|--|
| Target | EPHA10 |
| Synonyms | DFNA88 |
| Description | Recombinant human EPHA10 Protein with C-terminal 10×His tag |
| Delivery | In Stock |
| Uniprot ID | Q5JZY3 |
| Expression Host | HEK293 |
| Tag | C-10×His tag |
| Molecular Characterization | EPHA10(Glu34-Ala565) 10×His tag |
| Molecular Weight | The protein has a predicted molecular mass of 59.0 kDa after removal of the signal peptide. |
| Purity | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | Ephrin receptors, the largest subfamily of receptor tyrosine kinases (RTKs), and their ephrin ligands are important mediators of cell-cell communication regulating cell attachment, shape, and mobility in neuronal and epithelial cells (Aasheim et al., 2005 [PubMed 15777695]). See MIM 179610 for additional background on Eph receptors and ephrins.[supplied by OMIM, Mar 2008] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human EPHA10 Protein, His Tag on SDS-PAGE under reducing condition.

