

## **PRODUCT INFORMATION**

F2RL2 **Target** 

F2RL2, PAR3, Protease-activated receptor 3, **Synonyms** Thrombin receptor-like 2, Coagulation factor II

receptor-like 2

Recombinant human F2RL2 Protein with C-**Description** 

terminal human Fc tag

Delivery In Stock **Uniprot ID** 000254 **HEK293 Expression Host** 

Tag C-Human Fc tag

Molecular

Storage & Shipping

**Background** 

**Purity** 

F2RL2(Thr39-Thr94) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of Molecular Weight 32.2 kDa after removal of the signal peptide.

The purity of the protein is greater than 95% as

determined by SDS-PAGE and Coomassie blue

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

F2RL2 (Protease-activated receptor 3, PAR3) is a G-protein coupled receptor (GPCR) activated by thrombin and other proteases. It primarily couples to Gq/11 and Gi proteins, initiating phospholipase C activation, intracellular Ca<sup>2+</sup> mobilization, and MAPK signaling. F2RL2 is expressed in platelets, endothelial cells, and vascular smooth muscle, where it participates in hemostasis, platelet

activation, vascular inflammation, and thrombosis. Dysregulation of F2RL2 signaling is

linked to coagulation disorders and

cardiovascular diseases, making it a potential therapeutic target in thrombosis and vascular

pathology.

Usage Research use only

Conjugate Unconjugated

> Email: info@dimabio.com Website: www.dimabio.com





Human F2RL2 Protein, hFc Tag

Cat. No. PME101911



Figure 1. Human F2RL2 Protein, hFc Tag on SDS-PAGE under reducing condition.

15





