

PRODUCT INFORMATION

C-Flag&Strep Tag Tag

FPR2 **Target**

ALXR; FMLP-R-II; FMLPX; FPR2A; FPRH1; FPRH2; **Synonyms**

FPRL1; HM63; LXA4R

Human FPR2-Strep full length protein-synthetic Description

nanodisc

Delivery In Stock **Uniprot ID** P25090 **HFK293 Expression Host**

Formulation & Reconstitution

Protein Families Druggable Genome, GPCR, Transmembrane

Protein Pathways Neuroactive ligand-receptor interaction

The human full length FPR2-Strep protein has a **Molecular Weight**

MW of 39.0 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

Low affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophil chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system. The activation of LXA4R could result in an anti-inflammatory outcome

Background

counteracting the actions of proinflammatory signals such as LTB4 (leukotriene B4). Receptor for the chemokine-like protein FAM19A5, mediating FAM19A5-stimulated macrophage chemotaxis and the inhibitory effect on

> Email: info@dimabio.com Website: www.dimabio.com

TNFSF11/RANKL-induced osteoclast

differentiation.

Research use only Usage

Conjugate Unconjugated





ELISA assay to evaluate FPR2-Strep-Nanodisc 0.2µg Human FPR2-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag FPR2-Strep-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with FPR2-Strep-nanodisc is 2.281ng/ml.

Email: info@dimabio.com Website: www.dimabio.com




Figure 2. Human FPR2-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

