

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	FXYP7
Synonyms	N/A
Description	Human FXYP7-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P58549
Expression Host	HEK293
Protein Families	Ion Channels: Other
Protein Pathways	N/A
Molecular Weight	The human full length FXYP7-Strep protein has a MW of 8.5 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This reference sequence was derived from multiple replicate ESTs and validated by similar human genomic sequence. This gene encodes a member of a family of small membrane proteins that share a 35-amino acid signature sequence domain, beginning with the sequence PFXYP and containing 7 invariant and 6 highly conserved amino acids. The approved human gene nomenclature for the family is FXYP-domain containing ion transport regulator. Transmembrane topology has been established for two family members (FXYP1 and FXYP2), with the N-terminus extracellular and the C-terminus on the cytoplasmic side of the membrane. FXYP2, also known as the gamma subunit of the Na,K-ATPase, regulates the properties of that enzyme. FXYP1 (phospholemman), FXYP2 (gamma), FXYP3 (MAT-8), FXYP4 (CHIF), and FXYP5 (RIC) have been shown to induce channel activity in experimental expression systems. This gene product, FXYP7, is novel and has not been characterized as a protein. [RefSeq curation by Kathleen J. Sweadner, Ph.D., sweadner@helix.mgh.harvard.edu , Dec 2000]
Usage	Research use only
Conjugate	Unconjugated

