

## PRODUCT INFORMATION

|   |   |
|---|---|
| <b>Tag</b>                              | C-Flag&Strep Tag  |
| <b>Target</b>                           | FZD1  |
| <b>Synonyms</b>                         | N/A   |
| <b>Description</b>                      | Human FZD1-Strep full length protein-synthetic nanodisc   |
| <b>Delivery</b>                         | In Stock  |
| <b>Uniprot ID</b>                       | Q9UP38  |
| <b>Expression Host</b>                  | HEK293  |
| <b>Protein Families</b>                 | Druggable Genome, GPCR, Transmembrane   |
| <b>Protein Pathways</b>                 | Basal cell carcinoma, Colorectal cancer, Melanogenesis, Pathways in cancer, Wnt signaling pathway   |
| <b>Molecular Weight</b>                 | The human full length FZD1-Strep protein has a MW of 71.2 kDa   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Background</b>                       | Members of the 'frizzled' gene family encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The FZD1 protein contains a signal peptide, a cysteine-rich domain in the N-terminal extracellular region, 7 transmembrane domains, and a C-terminal PDZ domain-binding motif. The FZD1 transcript is expressed in various tissues. |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |




**ELISA assay to evaluate FZD1-Strep-Nanodisc**
0.2 $\mu$ g Human FZD1-Strep-Nanodisc per well

Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag FZD1-Strep-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with FZD1-Strep-nanodisc is 1.312ng/ml.



Figure 2. Human FZD1-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

