

PRODUCT INFORMATION

| | |
|---|---|
| Tag | C-Flag&Strep Tag |
| Target | GHSR |
| Synonyms | GHDP |
| Description | Human GHSR-Strep full length protein-synthetic nanodisc |
| Delivery | In Stock |
| Uniprot ID | Q92847 |
| Expression Host | HEK293 |
| Protein Families | Druggable Genome, GPCR, Transmembrane |
| Protein Pathways | Neuroactive ligand-receptor interaction |
| Molecular Weight | The human full length GHSR-Strep protein has a MW of 41.3 kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein may play a role in energy homeostasis and regulation of body weight. Two identified transcript variants are expressed in several tissues and are evolutionary conserved in fish and swine. One transcript, 1a, excises an intron and encodes the functional protein; this protein is the receptor for the Ghrelin ligand and defines a neuroendocrine pathway for growth hormone release. The second transcript (1b) retains the intron and does not function as a receptor for Ghrelin; however, it may function to attenuate activity of isoform 1a. Mutations in this gene are associated with autosomal idiopathic short stature. |
| Usage | Research use only |
| Conjugate | Unconjugated |



ELISA assay to evaluate GHSR-Strep-Nanodisc 0.2 μ g Human GHSR-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag GHSR-Strep-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with GHSR-Strep-nanodisc is 1.987ng/ml.

kDa M R



Figure 2. Human GHSR-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

