

PRODUCT INFORMATION

Target	GLP1R
Synonyms	GLP-1;GLP-1-R;GLP-1R
Description	Recombinant Human GLP1R Protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	P43220
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	GLP1R(Arg24-Glu139) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 39.7 kDa after removal of the signal peptide. The apparent molecular mass of GLP1R-hFc is approximately 35-70 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a 7-transmembrane protein that functions as a receptor for glucagon-like peptide 1 (GLP-1) hormone, which stimulates glucose-induced insulin secretion. This receptor, which functions at the cell surface, becomes internalized in response to GLP-1 and GLP-1 analogs, and it plays an important role in the signaling cascades leading to insulin secretion. It also displays neuroprotective effects in animal models. Polymorphisms in this gene are associated with diabetes. The protein is an important drug target for the treatment of type 2 diabetes and stroke. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Apr 2016]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human GLP1R Protein, hFc Tag on SDS-PAGE under reducing condition.

Human GLP1R, hFc Tagged Protein ELISA

0.2 μ g of Human GLP1R, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human GLP1R Protein, hFc Tag (PME101269) can bind Anti-GLP1R antibody(1A4), Rabbit mAb (DME101291) in a linear range of 0.32–1.6 ng/mL.

