

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	GPR132
Synonyms	G2A
Description	Human GPR132 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q9UNW8
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length GPR132 protein has a MW of 42.5 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Formulation & Reconstitution	The protein is a member of the guanine nucleotide-binding protein (G protein)-coupled receptor (GPCR) superfamily. The receptors are seven-pass transmembrane proteins that respond to extracellular cues and activate intracellular signal transduction pathways. This protein was reported to be a receptor for lysophosphatidylcholine action, but PubMedID: 15653487 retracts this finding and instead suggests this protein to be an effector of lysophosphatidylcholine action. This protein may have proton-sensing activity and may be a receptor for oxidized free fatty acids.
Storage&Shipping	Research use only
Background	Unconjugated
Usage	Research use only
Conjugate	Unconjugated




ELISA assay to evaluate GPR132-Nanodisc
0.2 μ g Human GPR132-Nanodisc per well

Figure 1. Elisa plates were pre-coated with Flag Tag GPR132-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with GPR132-Nanodisc is 4.100ng/ml.

kDa M R



Figure 2. WB analysis of Human GPR132-Nanodisc with anti-Flag monoclonal antibody at 1/5000 dilution, followed by Goat Anti-Rabbit IgG HRP at 1/5000 dilution

