

PRODUCT INFORMATION

Tag C-Flag&Strep Tag

Target GRIA3

Synonyms GLUR-C, GLUR-K3, GLUR3, GLURC, GluA3, MRX94,

MRXSW

DescriptionHuman GRIA3-Strep full length protein-synthetic

nanodisc 6~8weeks

Delivery 6∼8weeks
Uniprot ID P42263
Expression Host HEK293

Protein Families Ion Channels: Glutamate Receptors

Protein Pathways N/A

Background

Molecular Weight

The human full length GRIA3-Strep protein has a

MW of 101.2 kDa

Formulation & Reconstitution

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shippingat -80°C (Avoid repeated freezing and thawing).
Lyophilized proteins are shipped at ambient

temperature.

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. These receptors are heteromeric protein complexes composed of multiple subunits, arranged to form ligand-gated ion channels. The classification of glutamate receptors is based on their activation by different pharmacologic agonists. The subunit encoded by this gene belongs to a family of AMPA

encoded by this gene belongs to a family of AMPA (alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate)-sensitive glutamate receptors, and is subject to RNA editing (AGA->GGA; R->G). Alternative splicing at this locus results in different income which may be in their circular different income.

different isoforms, which may vary in their signal transduction properties. [provided by RefSeq, Jul

Email: info@dimabio.com Website: www.dimabio.com

2008]

Usage Research use only
Conjugate Unconjugated

