

## **PRODUCT INFORMATION**

Tag C-Flag Tag

**GRPR Target** 

**Synonyms** BB2; BB2R; BRS2

Human GRPR full length protein-synthetic **Description** 

nanodisc **Delivery** In Stock **Uniprot ID** P30550 **Expression Host HEK293** 

**Protein Families** Druggable Genome, GPCR, Transmembrane

Calcium signaling pathway, Neuroactive ligand-**Protein Pathways** 

receptor interaction

The human full length GRPR protein has a MW of **Molecular Weight** 

43.2 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before Formulation & Reconstitution lyophilization. Please see Certificate of Analysis

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

Store at -20°C to -80°C for 12 months in

temperature.

Gastrin-releasing peptide (GRP) regulates numerous functions of the gastrointestinal and central nervous systems, including release of gastrointestinal hormones, smooth muscle cell contraction, and epithelial cell proliferation and is a potent mitogen for neoplastic tissues. The effects of GRP are mediated through the gastrin-releasing peptide receptor. This receptor is a glycosylated, 7-transmembrane G-protein coupled receptor that activates the phospholipase

C signaling pathway. The receptor is aberrantly expressed in numerous cancers such as those of the lung, colon, and prostate. An individual with autism and multiple exostoses was found to have a balanced translocation between chromosome 8 and a chromosome X breakpoint located within the gastrin-releasing peptide receptor gene.

> Email: info@dimabio.com Website: www.dimabio.com

**Usage** Research use only Conjugate Unconjugated

**Background** 





## ELISA assay to evaluate GRPR-Nanodisc 0.2µg Human GRPR-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag GRPR-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with GRPR-Nanodisc is 4.434ng/ml.




Figure 2. Human GRPR-Nanodisc, Flag Tag on SDS-PAGE



Email: info@dimabio.com Website: www.dimabio.com

