

## PRODUCT INFORMATION

<b>Tag</b>	C-Flag Tag
<b>Target</b>	HCN1
<b>Synonyms</b>	BCNG-1, BCNG1, DEE24, EIEE24, GEFSP10, HAC-2
<b>Description</b>	Human HCN1 full length protein-synthetic nanodisc
<b>Delivery</b>	6~8weeks
<b>Uniprot ID</b>	060741
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Ion Channels: Cyclic nucleotide gated
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length HCN1 protein has a MW of 98.8kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The membrane protein encoded by this gene is a hyperpolarization-activated cation channel that contributes to the native pacemaker currents in heart and neurons. The encoded protein can homodimerize or heterodimerize with other pore-forming subunits to form a potassium channel. This channel may act as a receptor for sour tastes. [provided by RefSeq, Oct 2011]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated

