

## **PRODUCT INFORMATION**

**Target** Her2

**Synonyms** ERBB2;CD340;HER-2/neu;HER2;MLN19;NEU;NGL;TKR1

Recombinant human Her2 Protein with C-terminal Description

6×His tag In Stock **Delivery** 

**Uniprot ID** P04626 **Expression Host HEK293** 

Tag C-6×His Tag

Molecular

Reconstitution

**Background** 

Her2(Thr23-Thr652) 6×His tag Characterization

The protein has a predicted molecular mass of 70.2

kDa after removal of the signal peptide. The apparent molecular mass of Her2-His is approximately 100-130 **Molecular Weight** 

kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue **Purity** 

Formulation &

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for

specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid

Storage & Shipping repeated freezing and thawing). Lyophilized proteins

are shipped at ambient temperature.

This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinasemediated activation of downstream signalling

pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have

been reported, with the most common allele, lle654/lle655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding

different isoforms and others that have not been fully

Email: info@dimabio.com Website: www.dimabio.com

characterized. Usage Research use only Conjugate Unconjugated










Figure 1. Human Her2 Protein, His Tag on SDS-PAGE under reducing condition.

## Human Her2, His tagged protein ELISA

0.1 μg of Human Her2, His tagged protein per well




Figure 2. ELISA plate pre-coated by 1  $\mu$ g/ml (100  $\mu$ l/well) Human Her2,His tagged protein (PME100095) can bind Anti-Her2 Neutralizing antibody BME100048 in a linear range of 3.2-400 ng/ml.

Email: info@dimabio.com Website: www.dimabio.com

