

PRODUCT INFORMATION

IL-16 **Target**

Pro-Interleukin-16;Interleukin-16;IL-16;Lymphocyte **Synonyms**

Chemoattractant Factor; LCF; IL16

Recombinant Human Interleukin-16 is produced by Description our E.coli expression system and the target gene

encoding Met1-Ser130 is expressed.

Delivery In Stock

AAC12732.1 **Uniprot ID**

Expression Host E.coli

Molecular Weight

Background

Tag

Molecular Not available Characterization

Greater than 95% as determined by reducing SDS-**Purity**

PAGE.

13.4 KDa

Formulation & Lyophilized from a 0.2 µm filtered solution of

Reconstitution 20mM PB, 150mM NaCl, pH 7.0.

> Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Interleukin-16 (IL-16) is a CD8 T cell-derived cytokine that induces chemotaxis of CD4 T cells and CD4 monocytes and eosinophils. Analysis by gel filtration suggests that, under physiological conditions, human IL-16 exists predominantly as a noncovalently linked multimer, but that some IL-16 may exist as a monomer. However, only the multimeric form appears to possess chemotactic

activity, suggesting that receptor cross-linking may be required for activity. IL-16 also induces expression of IL-2 receptor (IL-2R) and MHC class II molecules on CD4 T cells. Human and murine IL-16 show significant cross-species reactivity.

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only Conjugate Unconjugated








Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

Email: info@dimabio.com Website: www.dimabio.com

