

## PRODUCT INFORMATION

<b>Target</b>	IL-29
<b>Synonyms</b>	Interferon lambda-1;IFN-lambda-1;Cytokine Zcyto21;Interleukin-29;IL-29;IFNL1;IL29;ZCYTO21
<b>Description</b>	Recombinant Human Interferon Lambda-1 is produced by our Mammalian expression system and the target gene encoding Gly20-Thr200 is expressed with a 10His tag at the C-terminus.
<b>Delivery</b>	Under development
<b>Uniprot ID</b>	Q8IU54
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-10×His Tag
<b>Molecular Characterization</b>	Not available
<b>Molecular Weight</b>	21.4 KDa
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Interleukin-29 (IL-29) is a secreted protein which belongs to the IL-28/IL-29 family. IL-29 is a cytokine with immunomodulatory activity. IL-29 is highly similar in amino acid sequence to the IL-28. IL-28 and IL-29 are induced by viral infection and showed antiviral activity. IL-28 and IL-29 interacted with a heterodimeric class II cytokine receptor that consisted of IL-10 receptor beta (IL-10R beta) and an orphan class II receptor chain, designated IL-28R alpha. IL-29 plays an important role in host defenses against microbes and its gene is highly upregulated in cells infected with viruses. IL-29 may play a role in antiviral immunity. IL-29 up-regulates MHC class I antigen expression. It is a Ligand for the heterodimeric class II cytokine receptor composed of IL10RB and IL28RA. The ligand/receptor complex seems to signal through the Jak-STAT pathway.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated

