

PRODUCT INFORMATION

Target	IL13
Synonyms	IL-13;NC30
Description	Recombinant human IL13(35-146) protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	P35225
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	IL13(Gly35-Asn146) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 38.4 kDa after removal of the signal peptide. The apparent molecular mass of IL13(35-146)-hFc is approximately 33-53 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen-induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human IL13(35-146) Protein, hFc Tag on SDS-PAGE under reducing condition.

