

PRODUCT INFORMATION

IL13RA1 **Target**

Synonyms CD213A1;CT19;IL-13Ra;NR4

Recombinant Human IL13RA1 Protein with C-**Description**

terminal 6×His tag

Delivery In Stock **Uniprot ID** P78552 **Expression Host HEK293** Tag C-6×His Tag

Molecular

Background

IL13RA1(Gly22-Thr343) 6×His tag Characterization

The protein has a predicted molecular mass of **Molecular Weight**

37.7 kDa after removal of the signal peptide. The apparent molecular mass of IL13RA1-His is approximately 55-70 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a subunit of the interleukin 13 receptor. This subunit forms a receptor complex with IL4 receptor alpha, a subunit shared by IL13 and IL4 receptors. This subunit serves as a primary IL13-binding subunit

of the IL13 receptor, and may also be a component of IL4 receptors. This protein has been shown to bind tyrosine kinase TYK2, and

thus may mediate the signaling processes that lead to the activation of JAK1, STAT3 and STAT6 induced by IL13 and IL4. [provided by RefSeq, Jul

Email: info@dimabio.com Website: www.dimabio.com

20081

Usage Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)








Figure 1. Human IL13RA1 Protein, His Tag on SDS-PAGE under reducing condition.

Human IL13RA1, His Tagged protein ELISA

0.2 µg of Human IL13RA1, HIs tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human IL13RA1 Protein, His Tag(PME100291) can bind Anti-IL13RA1 antibody(DM183), Rabbit mAb in a linear range of 0.32–8.0 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com



Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)



Human IL13RA1, His Tagged protein ELISA

0.2 μg of Human IL13RA1, His tagged protein per well




Figure 3. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human IL13RA1 Protein, His Tag (PME100291) can bind Anti-IL13RA1(eblasakimab biosimilar) mAb (BME100240) in a linear range of 0.64–80 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

