

PRODUCT INFORMATION

IL15 and IL15RA **Target Synonyms** IL-15,CD215

Recombinant human IL15(N120D) protein with C-**Description**

terminal human Fc tag and human IL15RA protein

Delivery

Uniprot ID Q13261 and P40933

Expression Host HEK293

Tag C-Human Fc tag and tag Free

Molecular IL15(Asn49-Ser162,N120D) hFc(Glu99-Ala330)

Characterization IL15RA(Ile31-Arg95)

The protein has a predicted molecular mass of 38.9kDa and 7.3kDa kDa after removal of the **Molecular Weight**

signal peptide.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue **Purity**

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before Reconstitution

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

Interleukin-15 is a member of the four α helix bundle family of cytokines. It plays a multifaceted role in the development and control of the immune system. The IL-15R consists of three polypeptides, the type-specific IL-15R α , the IL-2/IL-15R β , and the common γ -chain (γ C), which is shared by multiple cytokine receptors. IL-15 could enhance the effect of chemotherapeutic agents and ACT on transplanted tumors, but had

agents and ACT on transplanted tumors, but had little or no effect when administered alone. The result of research shows that the biological activity of IL-15 could be increased ~50-fold by administering preformed complexes of IL-15 and

its soluble receptor, IL-15Rα.

Usage Research use only

Conjugate Unconjugated

Background


Email: info@dimabio.com Website: www.dimabio.com





Human IL15RA & IL15, hFc Tagged (N120D) Protein ELISA

0.2 μg of Human IL15RA & IL15, hFc tagged protein per well



Anti-IL15(ordesekimab biosimilar) mAb (ng/mL)

Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human IL15RA & IL15 Protein, hFc Tag (N120D) (PME101891) can bind Anti-IL15(ordesekimab biosimilar) mAb (BME100253) in a linear range of 1.6–8 ng/mL.In order to specifically detect BME100253, mouse anti-human Fab-specific antibody was used as detection antibody.





Figure 1.Human IL15(N120D) protein with C-terminal human Fc tag and human IL15RA protein on SDS-PAGE under reducing condition.



Human IL15RA & IL15, hFc Tagged (N120D) Protein ELISA

0.2 μg of Human IL15RA & IL15, hFc tagged protein per well



Anti-IL15(ordesekimab biosimilar) mAb (ng/mL)

Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human IL15RA and IL15 Protein, hFc Tag (N120D) (PME101891) can bind Anti-IL15(ordesekimab biosimilar) mAb (BME100253) in a linear range of 1.6–8 ng/mL.In order to specifically detect BME100253, mouse anti-human Fab-specific antibody was used as detection antibody.

Email: info@dimabio.com Website: www.dimabio.com

