

PRODUCT INFORMATION

| | |
|------------------------------|--|
| Target | IL20RA |
| Synonyms | CRF2-8, IL-20R1, IL-20RA, IL-20R-alpha |
| Description | Recombinant human IL20RA Protein with C-terminal 10×His tag |
| Delivery | In Stock |
| Uniprot ID | Q9UHF4 |
| Expression Host | HEK293 |
| Tag | C-10×His tag |
| Molecular Characterization | IL20RA(Val30-Lys250) 10×His tag |
| Molecular Weight | The protein has a predicted molecular mass of 26.6 kDa after removal of the signal peptide. |
| Purity | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a member of the type II cytokine receptor family. The encoded protein is a subunit of the receptor for interleukin 20, a cytokine that may be involved in epidermal function. The interleukin 20 receptor is a heterodimeric complex consisting of the encoded protein and interleukin 20 receptor beta. This gene and interleukin 20 receptor beta are highly expressed in skin, and are upregulated in psoriasis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1.Human IL20RA Protein, His Tag on SDS-PAGE under reducing condition.

