

PRODUCT INFORMATION

Target IL20RA

CRF2-8, IL-20R1, IL-20RA, IL-20R-alpha **Synonyms**

Recombinant human IL20RA Protein with C-**Description**

terminal 10×His tag

Delivery In Stock **Uniprot ID** Q9UHF4 **Expression Host HEK293**

Tag C-10×His tag

Molecular

Purity

IL20RA(Val30-Lys250) 10×His tag Characterization

The protein has a predicted molecular mass of **Molecular Weight** 26.6 kDa after removal of the signal peptide.

The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before Formulation &

lyophilization. Please see Certificate of Analysis Reconstitution for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a member of the type II cytokine receptor family. The encoded protein is a subunit of the receptor for interleukin 20, a cytokine that may be involved in epidermal function. The interleukin 20 receptor is a


heterodimeric complex consisting of the encoded protein and interleukin 20 receptor beta. This **Background**

gene and interleukin 20 receptor beta are highly expressed in skin, and are upregulated in psoriasis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013]

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only

Conjugate Unconjugated



/+86-400-006-0995(China)








Figure 1.Human IL20RA Protein, His Tag on SDS-PAGE under reducing condition.

Email: info@dimabio.com Website: www.dimabio.com

