

PRODUCT INFORMATION

| | |
|------------------------------|---|
| Target | IL20RB |
| Synonyms | DIRS1; FNDC6; IL-20R2 |
| Description | Recombinant human IL20RB Protein with C-terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | Q6UXL0 |
| Expression Host | HEK293 |
| Tag | C-Human Fc tag |
| Molecular Characterization | IL20RB(Asp30-Leu233) hFc(Glu99-Ala330) |
| Molecular Weight | The protein has a predicted molecular mass of 48.9 kDa after removal of the signal peptide. |
| Purity | The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | IL20RB and IL20RA (MIM 605620) form a heterodimeric receptor for interleukin-20 (IL20; MIM 605619) (Blumberg et al., 2001 [PubMed 11163236]).[supplied by OMIM, Feb 2009] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human IL20RB Protein, hFc Tag on SDS-PAGE under reducing condition.

