

PRODUCT INFORMATION

| | |
|------------------------------|--|
| Target | IL36G |
| Synonyms | IL1E; IL1F9; IL1H1; IL-1F9; IL-1H1; IL1RP2; IL-1RP2 |
| Description | Recombinant human IL36G Protein with C-terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | Q9NZH8 |
| Expression Host | HEK293 |
| Tag | C-Human Fc tag |
| Molecular Characterization | IL36G(Ser18-Asp169) hFc(Glu99-Ala330) |
| Molecular Weight | The protein has a predicted molecular mass of 43.2 kDa after removal of the signal peptide. |
| Purity | The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein encoded by this gene is a member of the interleukin 1 cytokine family. The activity of this cytokine is mediated by interleukin 1 receptor-like 2 (IL1RL2/IL1R-rp2), and is specifically inhibited by interleukin 1 family, member 5 (IL1F5/IL-1 delta). Interferon-gamma, tumor necrosis factor-alpha and interleukin 1, beta (IL1B) are reported to stimulate the expression of this cytokine in keratinocytes. The expression of this cytokine in keratinocytes can also be induced by a contact hypersensitivity reaction or herpes simplex virus infection. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. [provided by RefSeq, May 2019] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human IL36G Protein, hFc Tag on SDS-PAGE under reducing condition.

