

PRODUCT INFORMATION

Target	IL5
Synonyms	IL-5;TRF;Interleukin-5;EDF
Description	Recombinant Human IL5 protein with C-terminal 6×His tag
Delivery	In Stock
Uniprot ID	P05113
Expression Host	HEK293
Tag	C-6×His Tag
Molecular Characterization	IL5(Ile20-Ser134) 6×His tag
Molecular Weight	The protein has a predicted molecular mass of 13.94 kDa after removal of the signal peptide.The apparent molecular mass of IL5-His is approximately 15-25 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. The encoded cytokine plays a major role in the regulation of eosinophil formation, maturation, recruitment and survival. The increased production of this cytokine may be related to pathogenesis of eosinophil-dependent inflammatory diseases. This cytokine functions by binding to its receptor, which is a heterodimer, whose beta subunit is shared with the receptors for interleukine 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene is located on chromosome 5 within a cytokine gene cluster which includes interleukin 4 (IL4), interleukin 13 (IL13), and CSF2 . This gene, IL4, and IL13 may be regulated coordinately by long-range regulatory elements spread over 120 kilobases on chromosome 5q31.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human IL5 Protein, His Tag on SDS-PAGE under reducing condition.




Figure 2. Flow cytometry analysis with 1  $\mu$ g/mL Human IL5 Protein, His Tag (PME100085) on HEK293 cells transfected with Human IL5RA protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).



**Human IL5, His tagged protein ELISA**  
0.5 µg of Human IL5, His tagged protein per well




Figure 3. ELISA plate pre-coated by 5 µg/mL (100 µL/well) Human IL5 protein, His Tag (PME100085) can bind Human IL5RA Protein, hFc Tag PME100536 in a linear range of 6.4-4000 ng/mL.

