

PRODUCT INFORMATION

| | |
|-----------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Tag | C-Flag&Strep Tag |
| Target | KCD12 |
| Synonyms | C13orf2, PFET1, PFETIN |
| Description | Human KCD12-Strep full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | Q96CX2 |
| Expression Host | HEK293 |
| Protein Families | Ion Channels: Other |
| Protein Pathways | N/A |
| Molecular Weight | The human full length KCD12-Strep protein has a MW of 35.7 kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | Auxiliary subunit of GABA-B receptors that determine the pharmacology and kinetics of the receptor response. Increases agonist potency and markedly alter the G-protein signaling of the receptors by accelerating onset and promoting desensitization (By similarity).[UniProtKB/Swiss-Prot Function] |
| Usage | Research use only |
| Conjugate | Unconjugated |

