

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	LPAR1
Synonyms	edg-2; EDG2; Gpcr26; LPA1; Mrec1.3; rec.1.3; vzg-1; VZG1
Description	Human LPAR1-Strep full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q92633
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Gap junction, Neuroactive ligand-receptor interaction
Molecular Weight	The human full length LPAR1-Strep protein has a MW of 41.1 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The integral membrane protein is a lysophosphatidic acid (LPA) receptor from a group known as EDG receptors. These receptors are members of the G protein-coupled receptor superfamily. Utilized by LPA for cell signaling, EDG receptors mediate diverse biologic functions, including proliferation, platelet aggregation, smooth muscle contraction, inhibition of neuroblastoma cell differentiation, chemotaxis, and tumor cell invasion.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate LPAR1-Strep-Nanodisc
0.2µg Human LPAR1-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag LPAR1-Strep-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with LPAR1-Strep-nanodisc is 8.022ng/ml.




Figure 2. Human LPAR1-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

