

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	LPAR2
Synonyms	EDG-4; EDG4; LPA-2; LPA2
Description	Human LPAR2-Strep full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q9HBW0
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Neuroactive ligand-receptor interaction
Molecular Weight	The human full length LPAR2-Strep protein has a MW of 38.7 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	A member of family I of the G protein-coupled receptors, as well as the EDG family of proteins. This protein functions as a lysophosphatidic acid (LPA) receptor and contributes to Ca ²⁺ mobilization, a critical cellular response to LPA in cells, through association with Gi and Gq proteins. An alternative splice variant has been described but its full length sequence has not been determined.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate LPAR2-Strep-Nanodisc
0.2 μ g Human LPAR2-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag LPAR2-Strep-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with LPAR2-Strep-nanodisc is 5.728ng/ml.



Figure 2. Human LPAR2-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

