

## **PRODUCT INFORMATION**

M-CSF **Target** 

Macrophage Colony-Stimulating Factor 1;CSF-1;M-**Synonyms** 

CSF;MCSF;Lanimostim;CSF1

Recombinant Human Macrophage Colony-

Stimulating Factor is produced by our Mammalian expression system and the target gene encoding Glu33-Arg255 is expressed with a 6His tag at the Description

C-terminus.

**Delivery** In Stock P09603 **Uniprot ID Expression Host HEK293** C-6×His Tag Tag

Molecular Not available Characterization

**Molecular Weight** 

Storage & Shipping

**Background** 

Greater than 95% as determined by reducing **Purity** 

26.17 KDa

Lyophilized from a 0.2  $\mu m$  filtered solution of 20mM PB, 150mM NaCl, pH 7.2. Formulation & Reconstitution

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

Macrophage Colony-Stimulating Factors (m-csf) are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and themonocytes-macrophages. CSF-1 promotes the release of proinflammatory chemokines, and thereby plays an important role in innate immunity and in inflammatory processes. It also plays an

important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone development. CSF-1 is required for normal male and female fertility and promotes reorganization of the actin system of of the actin cytoskeleton, regulates formation of membrane ruffles, cell adhesion and cell migration. it also plays a role in lipoprotein

> Email: info@dimabio.com Website: www.dimabio.com

clearance.

Usage Research use only Conjugate Unconjugated








Figure 1. Greater than 95% as determined by reducing SDS-PAGE.



Email: info@dimabio.com Website: www.dimabio.com