

PRODUCT INFORMATION

Target	MMP11
Synonyms	MMP-11;SL-3;ST3;STMY3
Description	Recombinant human MMP11 protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	P24347
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	MMP11(Arg32-Leu488) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 76.5 kDa after removal of the signal peptide. The apparent molecular mass of MMP11-hFc is approximately 70-100 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. However, the enzyme encoded by this gene is activated intracellularly by furin within the constitutive secretory pathway. Also in contrast to other MMP's, this enzyme cleaves alpha 1-proteinase inhibitor but weakly degrades structural proteins of the extracellular matrix. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human MMP11 Protein, hFc Tag on SDS-PAGE under reducing condition.

