

PRODUCT INFORMATION

MUC16 **Target Synonyms** CA125

Recombinant human MUC16(13810-14451) Description

Protein with C-terminal 10×His tag

Delivery In Stock **Uniprot ID** Q8WXI7 **Expression Host HEK293** Tag C-10×His tag

Storage & Shipping

Background

Molecular MUC16(Pro13810-Pro14451) 10×His tag Characterization

> The protein has a predicted molecular mass of 73.2 kDa after removal of the signal peptide. The

Molecular Weight apparent molecular mass of

MUC16(13810-14451)-His is approximately

100-130 kDa due to glycosylation.

The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a protein that is a member of the mucin family. Mucins are high molecular weight, O-glycosylated proteins that play an important role in forming a protective mucous barrier, and are found on the apical surfaces of the epithelia. The encoded protein is a membrane-tethered mucin that contains an extracellular domain at its amino terminus, a large tandem repeat domain, and a

transmembrane domain with a short cytoplasmic domain. The amino terminus is highly

glycosylated, while the repeat region contains 156 amino acid repeats unit that are rich in serines, threonines, and prolines. Interspersed within the repeats are Sea urchin sperm protein Enterokinase and Agrin (SEA) modules, leucine-rich repeats and ankyrin (ANK) repeats. These

regions together form the ectodomain, and there is a potential cleavage site found near an SEA module close to the transmembrane domain. This protein is thought to play a role in forming a barrier, protecting epithélial cells from pathogens. Products of this gene have been used as a marker for different cancers, with higher expression levels associated with poorer outcomes. [provided by DefSoz May 2017]

Email: info@dimabio.com

Website: www.dimabio.com

by RefSeq, May 2017]

Usage Research use only

Unconjugated Conjugate

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)






Figure 1. Human MUC16(13810-14451) Protein, His Tag on SDS-PAGE under reducing condition.

Email: info@dimabio.com Website: www.dimabio.com

