

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	NPSR1
Synonyms	ASRT2; GPR154; GPRA; NPSR; PGR14; VRR1
Description	Human NPSR1 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q6W5P4
Expression Host	HEK293
Protein Families	Druggable Genome, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length NPSR1 protein has a MW of 42.7 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	A member of the vasopressin/oxytocin subfamily of G protein-coupled receptors. The encoded membrane protein acts as a receptor for neuropeptide S and affects a variety of cellular processes through its signaling. Increased expression of this gene in ciliated cells of the respiratory epithelium and in bronchial smooth muscle cells is associated with asthma. Polymorphisms in this gene have also been associated with asthma susceptibility, panic disorders, inflammatory bowel disease, and rheumatoid arthritis. Alternative splicing results in multiple transcript variants.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate NPSR1-Nanodisc
0.2µg Human NPSR1-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag NPSR1-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with NPSR1-Nanodisc is 6.386ng/ml.




Figure2. Human NPSR1-Nanodisc, Flag Tag on SDS-PAGE

