

PRODUCT INFORMATION

Target NPTX2

Synonyms NP2; NARP; NP-II

Recombinant human NPTX2 Protein with C-terminal **Description**

10×His tag

Delivery In Stock P47972 **Uniprot ID Expression Host HEK293**

Tag C-10×His tag

Molecular

Background

NPTX2(Gly16-Leu431) 10×His tag Characterization

The protein has a predicted molecular mass of 47.0 Molecular Weight

kDa after removal of the signal peptide.

The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue **Purity**

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Formulation & Reconstitution

specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use Storage & Shipping within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins

are shipped at ambient temperature.

This gene encodes a member of the family of

This gene encodes a member of the family of neuronal petraxins, synaptic proteins that are related to C-reactive protein. This protein is involved in excitatory synapse formation. It also plays a role in clustering of alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA)-type glutamate receptors at established synapses, resulting in non-apoptotic cell death of dopaminergic nerve cells. Upregulation of this gene in Parkinson disease (PD) tissues suggests that the protein may be involved in the pathology of PD. [provided by RefSeq, Feb 2009]

Usage Research use only Conjugate Unconjugated










Figure 1. Human NPTX2 Protein, His Tag on SDS-PAGE under reducing condition.

