

## PRODUCT INFORMATION

<b>Target</b>	NPY2R
<b>Synonyms</b>	NPY2-R
<b>Description</b>	Recombinant human NPY2R Protein with C-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P49146
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-Human Fc tag
<b>Molecular Characterization</b>	NPY2R(Met1-Val51) hFc(Glu99-Ala330)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 31.7 kDa after removal of the signal peptide.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Predicted to enable calcium channel regulator activity and neuropeptide Y receptor activity. Involved in cardiac left ventricle morphogenesis and outflow tract morphogenesis. Located in cilium. Implicated in Huntington's disease; morbid obesity; and obesity. Biomarker of peripheral artery disease and temporal lobe epilepsy. [provided by Alliance of Genome Resources, Jun 2025]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Human NPY2R Protein, hFc Tag on SDS-PAGE under reducing condition.

