

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	NTSR2
Synonyms	NTR2
Description	Human NTSR2-Strep full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	O95665
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Neuroactive ligand-receptor interaction
Molecular Weight	The human full length NTSR2-Strep protein has a MW of 45.4 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural agonist for the receptor may exist.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate NTSR2-Strep-Nanodisc
0.2µg Human NTSR2-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag NTSR2-Strep-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with NTSR2-Strep-nanodisc is 2.690ng/ml.




Figure 2. Human NTSR2-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

