

PRODUCT INFORMATION

Target	OPRD1
Synonyms	DOP; DOR; DOR1; OPRD
Description	Recombinant human OPRD1 Protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	P41143
Expression Host	HEK293
Tag	C-Human Fc tag
Molecular Characterization	OPRD1(Met1-Ala47) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 30.5 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Enables G protein-coupled enkephalin receptor activity. Involved in several processes, including G protein-coupled opioid receptor signaling pathway; cellular response to hypoxia; and regulation of mitochondrial membrane potential. Located in plasma membrane. Implicated in alcohol dependence; alcohol use disorder; drug dependence (multiple); opioid abuse; and withdrawal disorder. Biomarker of opioid abuse. [provided by Alliance of Genome Resources, Jun 2025]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human OPRD1 Protein, hFc Tag on SDS-PAGE under reducing condition.

